

- 25 -

- weight. Yet other advantages arise at 45 to 65% internal phase volume. Compositions with high proportion of disperse phase, i.e. from 65 to 85% disperse phase, may also be advantageous. They can give good hardness even though the concentration of esterified saccharide structurant may be only a small percentage of the total composition.

- An emulsion composition will generally include one or more emulsifying surfactants which may be anionic, cationic, zwitterionic and/or nonionic surfactants. The proportion of emulsifier in the composition is often selected in the range up to 10% by weight and in many instances from 0.1 or 0.25 up to 5% by weight of the composition. Most preferred is an amount from 0.1 or 0.25 up to 2 or 3% by weight, such as 0.3%, 0.4% or 0.5% by weight, or an amount in between. Nonionic emulsifiers are frequently classified by HLB value. It is desirable to use an emulsifier or a mixture of emulsifiers with an overall HLB value in a range from 2 to 10 preferably from 3 to 8.

- It may be convenient to use a combination of two or more emulsifiers which have different HLB values above and below the desired value. By employing the two emulsifiers together in appropriate ratio, it is readily feasible to attain a weighted average HLB value that promotes the formation of an emulsion.

- Many suitable emulsifiers of high HLB are nonionic ester or ether emulsifiers comprising a polyoxyalkylene moiety, especially a polyoxyethylene moiety, often containing from about 2 to 80, and especially 5 to 60 oxyethylene units,

- and/or contain a polyhydroxy compound such as glycerol or sorbitol or other alditol as hydrophilic moiety. The hydrophilic moiety can contain polyoxypropylene. The emulsifiers additionally contain a hydrophobic alkyl, alkenyl or aralkyl moiety, normally containing from about 8 to 50 carbons and particularly from 10 to 30 carbons. The hydrophobic moiety can be either linear or branched and is often saturated, though it can be unsaturated, and is optionally fluorinated. The hydrophobic moiety can comprise a mixture of chain lengths, for example those deriving from tallow, lard, palm oil, sunflower seed oil or soya bean oil. Such nonionic surfactants can also be derived from a polyhydroxy compound such as glycerol or sorbitol or other alditols. Examples of emulsifiers include cetareth-10 to 25, ceteth-10-25, steareth-10-25 (i.e.  $C_{16}$  to  $C_{18}$  alcohols ethoxylated with 10 to 25 ethylene oxide residues) and PEG-15-25 stearate or distearate. Other suitable examples include  $C_{10}$ - $C_{20}$  fatty acid mono, di or tri-glycerides. Further examples include  $C_{18}$ - $C_{22}$  fatty alcohol ethers of polyethylene oxides (8 to 12 EO).

- Examples of emulsifiers, which typically have a low HLB value, often a value from 2 to 6 are fatty acid mono or possibly diesters of polyhydric alcohols such as glycerol, sorbitol, erythritol or trimethylolpropane. The fatty acyl moiety is often from  $C_{14}$  to  $C_{22}$  and is saturated in many instances, including cetyl, stearyl, arachidyl and behenyl. Examples include monoglycerides of palmitic or stearic acid, sorbitol mono or diesters of myristic, palmitic or stearic acid, and trimethylolpropane monoesters of stearic acid.

0982077-104704

A particularly desirable class of emulsifiers comprises dimethicone copolymers, namely polyoxyalkylene modified dimethylpolysiloxanes. The polyoxyalkylene group is often a polyoxyethylene (POE) or polyoxypropylene (POP) or a copolymer of POE and POP. The copolymers often terminate in C<sub>1</sub> to C<sub>12</sub> alkyl groups. An especially desirable example of this class is available under the trade name ABIL EM90 for use within the aforementioned ranges of proportions.

10 Suitable emulsifiers and co-emulsifiers are widely available under many trade names and designations including Abil™, Arlacel™, Brij™, Cremophor™, Dehydrol™, Dehymuls™, Emerest™, Lameform™, Pluronic™, Prisorine™, Quest PGPR™, Span™, Tween™, SF1228, DC3225C and Q2-5200.

15

#### Antiperspirant Actives

If the composition is an antiperspirant, it will contain an antiperspirant active. Antiperspirant actives, are preferably incorporated in an amount of from 0.5-60%, particularly from 5 to 30% or 40% and especially from 5 or 10% to 30 or 35% of the weight of the composition.

Antiperspirant actives for use herein are often selected from astringent active salts, including in particular aluminium, zirconium and mixed aluminium/zirconium salts, including both inorganic salts, salts with organic anions and complexes. Preferred astringent salts include aluminium, zirconium and aluminium/zirconium halides and halohydrate salts, such as chlorohydrates.

25  
30